Atti dell'Accademia Peloritana dei Pericolanti Classe di Scienze Fisiche, Matematiche e Naturali Vol. LXXXVI, C1S0801005 (2008) - Suppl. 1

SUPERPOSITIONS IN PRIGOGINE'S APPROACH TO IRREVERSIBILITY FOR PHYSICAL AND FINANCIAL APPLICATIONS

DAVID CARFÌ

ABSTRACT. In this paper we apply the theory of superpositions for Radon measures on compact subsets of the real euclidean *n*-space \mathbb{R}^n to Prigogine's approach in the study of irreversible processes, which emerge in Physics and in Economics; showing that the superposition is a natural rigorous tool feasible to face the problem.

1. Introduction

In section 2 we give a definition of discrete dynamical system that will be generalized in the paper. In section 3 we expose a particularly important kind of discrete dynamical systems: the systems generated by a function from a set to itself; we show their basic properties. In section 4 we define the probabilistic dynamical systems and explain the necessity of a generalization of the Prigogine's intuitive settings. In section 5, 6 and 8 we develop the foundation of superpositions theory on compact subsets and on locally compact subspaces of the real euclidean *n*-space \mathbb{R}^n . In section 7 we use superpositions to define the probabilistic system generated by a map. In section 9 we prove the main result of the paper: a generalization of one Prigogine's result. For what concerns the origin of the paper, it can be found in [1]-[6], for the theory of Radon measures see [7] and [8].

2. Discrete dynamical systems

Definition (of abstract discrete dynamical system). A discrete (non-reversible) dynamical system on a non-empty set X is an action of the additive monoid of natural numbers on the set X; in other terms, it is a function $s : \mathbb{N}_0 \times X \to X$ such that $s(0, x_0) = x_0$, for every $x_0 \in X$ and such that $s(m + n, x_0) = s(m, s(n, x_0))$, for every pair of natural numbers (m, n).

Terminology. Let s be a discrete dynamical system on X. The set X is called the space of states or the space of phases of the system s. Every element x of X is called a state of the system s. Every natural number n is called a time of the system s.

Remark. An action a of a monoid on a set X induces on the set a preorder in a natural way: a state x is said to precede a state y if there is an element m of the monoid that sends

x into y, i.e, such that a(m, x) = y. In general it is possible to find not-comparable states. If the monoid is a group then this preorder is an equivalence relation on the set.

Definition (of orbit). For every time $n \in \mathbb{N}$ and for every state $x_0 \in X$, the state $s(n, x_0)$ is called the state of s at the time n determined by the initial condition (state) x_0 . Fixed the state x_0 , the sequence $(s(n, x_0))_{n \in \mathbb{N}}$ is called the orbit or trajectory of the system starting from the initial state x_0 .

Definition (of equilibria). A state x_e is said an equilibrium-state for the system s if the orbit starting from x_e is constant of value x_e , that is $s(n, x_e) = x_e$, for every time n. If τ is a topology on the state space X, a state x_e is said a τ -asintotic equilibrium state for s under the initial condition x_0 if the orbit starting from $x_0 \tau$ -converges to x_e :

$$\tau \lim_{n \to +\infty} s(n, x_0) = x_e$$

Remark. Clearly, an equilibrium state is a τ -asintotic equilibrium, for every topology τ on the set X.

Interpretation. A discrete dynamical system is the model for physical systems whose state depends on time, considered as a discrete quantity, and on its initial condition (state), that is the state at time 0.

Let us explain the definitions with an elementary example.

Example. Let $X = \mathbb{R}$ be the space of states of the dynamical system

$$s: \mathbb{N}_0 \times \mathbb{R} \to \mathbb{R}: s(n, x_0) = x_0^{n+1}$$

Let us consider the orbits of the system starting from different initial conditions:

1) if $|x_0| < 1$, we have $\lim_{n \to +\infty} s(n, x_0) = 0$. The orbit starting from x_0 converges to the state 0, thus, with Prigogine, we say that the system s converges as intotically to the equilibrium-state $x_e = 0$ starting from state x_0 ;

2) if $x_0 > 1$, we have $\lim_{n \to +\infty} s(n, x_0) = +\infty$, in this case the orbit $s(\cdot, x_0)$ is divergent (in the sense of real sequences);

3) if $x_0 \leq -1$, the orbit $(s(n, x_0))_{n \in \mathbb{N}}$ is oscillating (in the sense of real sequences), and moreover, if $x_0 < -1$, it is absolutely divergent $\lim_{n \to +\infty} |s(n, x_0)| = +\infty$;

4) if $x_0 = 0$, we have $s(n, x_0) = 0$, for every $n \in \mathbb{N}_0$; thus the state $x_0 = 0$ is an equilibrium state of the system.

5) if $x_0 = 1$, we have $s(n, x_0) = 1$, for every $n \in \mathbb{N}_0$, thus also $x_0 = 1$ is an equilibrium state of the system s.

Remark. Let a be an action of a monoid on a set X. We recall that a point e of a preordered space is said strongly maximal if the relation "x is weakly greater than e" implies that x is equal to e. It is clear that a point e of the set X is an equilibrium of the dynamical system if and only if it is a strongly maximal point of X with respect to the preorder induced by the action on the set. An element x of the set X is said a torsion element with respect to the action if there is an element of the monoid that sends x into itself. The class of indifference of a torsion element coincides with its orbit and an element

is a torsion element with respect to the action if and only if it is a maximal element with respect to the preorder induced by the action (for the proofs and a complete treatment see [9]).

3. Discrete dynamical systems generated by maps.

Definition (of discrete dynamical system generated by a map). Let X be a nonempty set (that will be the space of states of our dynamical system) and let $f : X \to X$ a function. Then we consider the mapping

$$s_f: \mathbb{N}_0 \times X \to X: (n, x_0) \mapsto f^n(x_0),$$

where by f^n we shall denote the composition of f with f itself n times. The mapping s_f is clearly a discrete dynamical system and it is called the system generated by the map f.

The equilibrium-states of a dynamical system generated by a map f are connected with the fixed point of f, as it is shown by the following theorems.

Theorem. Let $f : X \to X$ be a map. Then, i) every fixed point x_0 of f is an equilibrium point associated with x_0 ; ii) every fixed point of f in an orbit $s_f(\cdot, x)$ is an equilibrium associated with x; iii) every equilibrium of s_f is a fixed point of f; iv) if $s_f(n, x) = s_f(n + 1, x)$ for some n and $x, s_f(n, x)$ is an equilibrium under x.

Proof. i) If $x_0 \in Fix(f)$, we have $s_f(n, x_0) = f^n(x_0) = x_0$; hence x_0 is an equilibrium-state of the system s associated with the initial state x_0 , and more the orbit starting from x_0 has a single state.

ii) If for some $m \in \mathbb{N}_0$, we have $s_f(m, x) \in \operatorname{Fix}(f)$, for every $n \ge m$, we have

$$s_f(n,x) = s_f(m,x),$$

and then $s_f(m, x)$ is an equilibrium of s_f associated with the initial state x.

iii) If $\overline{x} = s_f(m, x_0)$ is an equilibrium-state associated with the initial condition x_0 , we have $s_f(m+1, x_0) = s_f(m, x_0) = \overline{x}$, and taking into account that

$$s_f(m+1, x_0) = f(s_f(m, x_0)) = f(\overline{x})$$

we conclude $\overline{x} \in Fix(f)$.

iv) We have

$$f(s_f(m, x_0)) = s_f(n+1, x_0) = s_f(n, x_0),$$

so $s_f(m, x_0)$ is a fixed point of f belonging to the orbit $s_f(\cdot, x_0)$, and then by (ii) an equilibrium associated with x_0 .

Theorem. Let (X, τ) be a topological space and let $f : X \to X$ be a τ -continuous function. Let s_f be the discrete dynamical system generated by f, let x_0 be a state such that the orbit $s_f(\cdot, x_0)$ converges and let

$$L := \lim_{n \to +\infty} s_f(n, x_0) \,.$$

Then, the asintotic equilibrium L of the system s_f is a fixed point of f.

Proof. We have

$$f(L) = f\left(\lim_{n \to +\infty} s_f(n, x_0)\right) = \lim_{n \to +\infty} f\left(s_f(n, x_0)\right) =$$
$$= \lim_{n \to +\infty} f\left(f^n(x_0)\right) = \lim_{n \to +\infty} f^{n+1}(x_0) =$$
$$= \lim_{n \to +\infty} s_f(n+1, x_0) = L.$$

As we desire.

Theorem. Let (X, d) be a complete metric space and let $f : X \to X$ be a contraction on (X, d). Let s_f be the discrete dynamical system generated by f. Then, there exists only an equilibrium state L for s_f and, for every state x_0 , the orbit $s_f(\cdot, x_0)$ converges to L, that is the only fixed point of f.

Proof. It follows from the Banach fixed-point theorem.

We can introduce the main example of the paper.

Definition (of Bernoulli shift). We define (deterministic) Bernoulli shift the system

$$s_f: \mathbb{N}_0 \times \mathbb{R} \to \mathbb{R}: (n, x_0) \mapsto f^n(x_0),$$

generated by the map

$$f:[0,1] \to [0,1]: x \mapsto \begin{cases} 2x & \text{if } x \in [0,1/2[\\ 2x-1 & \text{if } x \in [1/2,1] \end{cases}.$$

4. Prigogine probabilistic systems and superpositions

Our aim, following Prigogine, is to associate with the Bernoully shift s_f , and more generally with a discrete dynamical system, a probabilistic dynamical system, i.e., a dynamical system whose space of states is a space of probability measures. To this end we must define rigorously the concept of probabilistic discrete dynamical system on a compact subset of \mathbb{R}^n . Our definition will be extremely natural. In the following we shall denote by P(K) the set of probability measure on a compact subset K of \mathbb{R}^n , that is the subset of the dual of the Banach space $(C^0(K), \|\cdot\|_{\infty})$ containing those non-negative functionals p whose value on the constant unitary real function 1_K is 1.

Definition (of probabilistic discrete dynamical system). A probabilistic discrete dynamical system on a compact subset K of \mathbb{R}^n is an action of the additive monoid of the natural numbers on the space of the Radon probability measures on the compact, in other terms, it is a discrete dynamical system on the space of the Radon probability measures on the compact; explicitly, it is a function $s : \mathbb{N}_0 \times P(K) \to P(K)$ such that $s(0, p_0) = p_0$, for every $p_0 \in P(K)$ and such that $s(m + n, p_0) = s(m, s(n, p_0))$, for every pair of natural numbers (m, n).

We desire the classic dynamical systems on K correspond to particular probabilistic systems; to this aim note that, for every x_0 belonging to K, the Dirac measure δ_{x_0} means

the certainty to find the system in the state x_0 . For example, if s_f is the system generated by a function $f : [0,1] \rightarrow [0,1]$, the condition that the system s_f is in the state $f(x_0)$ at time 1 is $s_f(1, x_0) = f(x_0)$, this can be expressed, according to Prigogine, by the "integral" equality

$$\delta(x - f(x_0)) = \int_0^1 \delta(y - x_0) \,\delta(x - f(y)) \, dy, \quad (1)$$

unfortunately this expression does not have a precise mathematical sense, we shall give it a sense by the concept of superposition of a family of measures defined on a compact subset of \mathbb{R}^n . Moreover, the Prigogine's idea is the following: generalizing, if at a time $n \in \mathbb{N}_0$ we have a probability distribution of states p_n on [0, 1], at the time n + 1 the system shall be in the probabilistic state p_{n+1} given by

$$U(p_n)(x) := p_{n+1}(x) = \int_0^1 p_n(y)\delta(x - f(y)) \, dy.$$

regrettably, this expression holds, in the sense of distributions, when p_n and p_{n+1} are functions (and not too strange), but so we cannot use this expression for not-regular measures, and in particular we cannot generalize the equality (1). A correct and not-restrictive way to solve the problem can be found in the following section.

5. Superpositions of C^0 -family in the space $C^{0'}(K)$

In the following, if H is a compact subset of \mathbb{R}^n , we denote by $C^{0'}(H)$ the dual of the Banach space $(C^0(H), \|\cdot\|_{\infty})$.

Definition (families of class C^0). Let $K \subseteq \mathbb{R}^n$ and $H \subseteq \mathbb{R}^m$ be compact subsets. Let $v = (v_i)_{i \in H}$ be a family in $C^{0'}(K)$ indexed by H. We define image of a continuous test function ϕ on K by the family v, the function

$$v(\phi): H \to \mathbb{R}: v(\phi)(i) = v_i(\phi).$$

If the image $v(\phi)$ lies in $C^0(H)$, for every $\phi \in C^0(K)$, we say that v is a family of class C^0 . In these conditions we call the operator

$$\widehat{v}: C^0(K) \to C^0(H): \phi \mapsto v(\phi)$$

operator generated by v.

Definition (superposition of measures). Let $K \subseteq \mathbb{R}^n$ and $H \subseteq \mathbb{R}^m$ be compact subsets. Let $a \in C^{0'}(H)$, i.e., let a be a Radon-measure on H and let v be a family of class C^0 in $C^{0'}(K)$. The functional $a \circ \hat{v}$ is denoted by

$$\int_{H} av,$$

and is called superposition of v by the system of coefficients a. Moreover, if a is a probabilistic system of coefficients, i.e., if a is non-negative and with unitary integral, the superposition $\int_{H} av$ is called a convex superposition of v.

Theorem. Let $K \subseteq \mathbb{R}^n$ and $H \subseteq \mathbb{R}^m$ be compact subsets. Let $a \in C^{0'}(H)$, and let v be a family of class C^0 in the space $C^{0'}(K)$ indexed by the compact H. Then the functional

$$\int_{H} av : C^{0}(K) \to \mathbb{R},$$

is a Radon-measure, i.e., it is a continuous linear functional on $C^0(K)$. Moreover, if v is a family of probabilistic Radon-measures, every convex superposition of v is a probabilistic Radon-measure too.

Proof. Let ϕ be a continuous test function on K, we have, for some $j \in H$,

$$\left| \left(\int_{H} av \right) (\phi) \right| = |a (v (\phi))| \le ||a||_{C^{0'}(H)} ||v (\phi)||_{C^{0}(H)} = ||a||_{C^{0'}(H)} \max_{i \in H} |v_{i} (\phi)| = ||a||_{C^{0'}(H)} ||v_{j}||_{C^{0'}(K)} ||\phi||_{C^{0}(K)},$$

so $\int_{H} av$ is a bounded-linear functional and then it is continuous.

Let $v = (v_i)_{i \in H}$ be a C^0 -family of probability measures on K, i.e., let v be a family of Radon-measures on K such that every v_i is normalized and non-negative, i.e., v_i $(1_K) = 1$ for every $i \in H$, and v_i $(\phi) \ge 0$, for every $\phi \in C^0$ (K) such that $\phi \ge 0$. Let $a \in C^{0'}(H)$ be (analogously) such that $a(1_H) = 1$ and $a \ge 0$.

Normalization. We have

$$\left(\int_{H} av\right)(1_{K}) = a\left(v\left(1_{K}\right)\right);$$

now, for every i in H, we see $v(1_K)(i) = v_i(1_K) = 1$, that is equivalent to $v(1_K) = 1_H$, thus

$$\left(\int_{H} av\right)(1_{K}) = a\left(1_{H}\right) = 1.$$

Non-negativity. We have

$$\left(\int_{H} av\right)(\phi) = a\left(v\left(\phi\right)\right);$$

now $v(\phi)(i) = v_i(\phi) \ge 0$, for every $i \in H$, by non-negativity of v_i , hence $v(\phi)$ is a non negative function, and then by non-negativity of a, we desume $a(v(\phi)) \ge 0$, hence every C^0 -convex superposition of v is a probability measure.

Theorem. Let $f : H \to \mathbb{R}$ be a continuous function, let K be the range of f and let δ_f be the family $(\delta_{f(x)})_{x \in H}$ in the space $C^{0'}(K)$. Then, δ_f is a continuous family; moreover, for every measure $c \in C^{0'}(H)$, the superposition $\int_H c\delta_f$ is a measure in $C^{0'}(K)$ and we have

$$\left(\int_{H} c\delta_{f}\right)(\phi) = \int_{H} \phi \circ f \ d\mu_{c}.$$

Proof. Let us compute the image $\delta_f(\phi)$ for every $\phi \in C^0(K)$; we have

$$\delta_f(\phi): H \to \mathbb{R}, \ \delta_f(\phi)(x) = \delta_{f(x)}(\phi) = \phi(f(x)) = (\phi \circ f)(x),$$

hence $\delta_f(\phi) = \phi \circ f$; since $\phi \circ f \in C^0(H)$, δ_f is a C^0 -family. Let $c \in C^{0'}(H)$, we have so

$$\left(\int_{H} c\delta_{f}\right)(\phi) = c\left(\phi \circ f\right). \blacksquare$$

6. Superpositions of measurable-families in the space $C^{0'}(K)$

We desire to extend the definition of superposition to families v indexed by compact intervalls of the real line and piecewise-continuous. A function $f : [a, b] \rightarrow \mathbb{R}$ is said piecewise-continuous if it is continuous but in a finite set of points in which the right and left limits there exist finite and in which f is right or left continuous. To this end we shall define a larger collection of families: the class of measurable families.

Definition (measurable families). Let $K \subseteq \mathbb{R}^n$ and $H \subseteq \mathbb{R}^m$ be compact subsets and let $v = (v_i)_{i \in H}$ be a family in $C^{0'}(K)$ indexed by H. The image of a continuous test function ϕ on K under the family v, is again the function

$$v(\phi): H \to \mathbb{R}: v(\phi)(i) = v_i(\phi).$$

If the image of $v(\phi)$ is measurable in the sense of Borel (retro-image of open sets are Borel-measurable sets), for every $\phi \in C^0(K)$, we say that v is a measurable family. In these conditions, if $\mathcal{B}(H,\mathbb{R})$ is the set of Borel-measurable real functions on H, we call the operator

$$\widehat{v}: C^{0}(K) \to \mathcal{B}(H, \mathbb{R}): \phi \mapsto v(\phi),$$

operator generated by v.

If v is a measurable family and if $a \in C^{0'}(H)$, then the functional

$$\int_{H} av : C^{0}(K) \to \mathbb{R} : \left(\int_{H} av\right)(\phi) := \int_{H} v(\phi) \ d\mu_{a}$$

where μ_a is the unique Borel measure associated with the functional a by the Riesz Representation Theorem, is called superposition of v under the system of coefficients a.

Theorem. Let $a \in C^{0'}(H)$, and let v be a measurable family in the space $C^{0'}(K)$ indexed by the compact H. Then, if for every $\phi \in C^0(K)$ the function $|v(\phi)|$ attains his maximum, the functional

$$\int_{H} av : C^{0}(K) \to \mathbb{R},$$

is a mensure-distribution, i.e., a continuous linear functional on $C^0(K)$. Moreover, if v is a family of probabilistic mensural distributions, every convex superposition of v is a probabilistic mensural distribution too.

Proof. Let ϕ be a continuous test function on K, we have, for some $j \in H$,

$$\begin{split} \left| \left(\int_{H} av \right) (\phi) \right| &= \left| \int_{H} v(\phi) \, d\mu_{a} \right| \leq \int_{H} |v(\phi)| \, d \, |\mu_{a}| \leq |\mu_{a}| \, (H) \sup |v(\phi)| = \\ &= \|a\|_{C^{0'}(H)} \max_{i \in H} |v_{i}(\phi)| = \|a\|_{C^{0'}(H)} \, |v_{j}(\phi)| \leq \\ &\leq \|a\|_{C^{0'}(H)} \, \|v_{j}\|_{C^{0'}(K)} \, \|\phi\|_{C^{0}(K)} \,, \end{split}$$

so $\int_{H} av$ is a bounded-linear functional and then it is continuous. Let $v = (v_i)_{i \in H}$ be a C^0 -family of probability distributions on K, i.e., be such that every v_i is normalized and non-negative, i.e., $v_i (1_K) = 1$, for every $i \in H$, and $v_i (\phi) \ge 0$, for every $\phi \in C^0(K)$ such that $\phi \geq 0$. Let $a \in C^{0'}(H)$ be (analogously) such that $a(1_H) = 1$ and a > 0.

Normalization. We have

$$\left(\int_{H} av\right)(1_{K}) = \int_{H} v(1_{K}) d\mu_{a};$$

now, for every i in H, we see $v(1_K)(i) = v_i(1_K) = 1$, that is equivalent to $v(1_K) = 1_H$, thus $\langle r \rangle \langle r \rangle$

$$\left(\int_{H} av\right)(1_{K}) = \int_{H} 1_{H} d\mu_{a} = a(1_{H}) = 1.$$

Non-negativity. We have

$$\left(\int_{H} av\right)(\phi) = \int_{H} v(\phi) \, d\mu_{a};$$

now $v(\phi)(i) = v_i(\phi) \ge 0$, for every $i \in H$, by non-negativity of v_i , hence $v(\phi)$ is a non negative function, and then by non-negativity of a,

$$\int_{H} v\left(\phi\right) d\mu_a \ge 0$$

hence every C^0 -convex superposition of v is a probability distribution.

Theorem. Let $f : H \to \mathbb{R}$ be a Borel-measurable function with compact range K and let δ_f be the family $(\delta_{f(x)})_{x\in H}$ in the space $C^{0'}(K)$. Then, δ_f is a measurable family, moreover, for every $c \in C^{0'}(H)$ the superposition $\int_{H} c\delta_{f}$ is a measure and in particular we have

$$\left(\int_{H} c\delta_{f}\right)(\phi) = \int_{H} \phi \circ f \ d\mu_{c}$$

Proof. For every $\phi \in C^0(K)$, we have

$$\delta_f(\phi): H \to \mathbb{R}, \ \delta_f(\phi) = \phi \circ f \in \mathcal{B}(H, \mathbb{R})$$

where $\mathcal{B}(H,\mathbb{R})$ is the set of Borel-measurable real functions on H. Thus δ_f is a measurablefamily. Note that since $|\phi|$ is continuous it attains its maximum on the compact f (H), and then $|\phi \circ f|$ attains its maximum on H. Concluding, by the preceding theorem, for every $c \in C^{0'}(H)$, the superposition $\int_{H} c\delta_{f}$ is a measure and we have

$$\left(\int_{H}c\delta_{f}\right)(\phi)=\int_{H}\phi\circ f\;d\mu_{c}.\blacksquare$$

7. Probabilistic systems via superpositions

Recall that our aim is to associate with the classic dynamical systems on a compact subset K probabilistic dynamical systems on the same subset. In particular we desire to associate with the dynamical system generated by a mapping $f: K \to K$ a probabilistic dynamical system on K. For instance, as we said, if s_f is the system generated by a function $f : [0,1] \rightarrow [0,1]$, the condition that the system s_f can be found in the state $f(x_0)$ at time 1 is $s_f(1, x_0) = f(x_0)$, this can be expressed, at last rigorously, in terms of superposition

$$\delta_{f(x_0)} = \int_{[0,1]} \delta_{x_0} \left(\delta_{f(y)} \right)_{y \in [0,1]},$$

we can read: the certainty to find the system s_f in the state $f(x_0)$ at time 1 is the superposition of the family $(\delta_{f(y)})_{y \in [0,1]}$ with respect to the system of coefficients δ_{x_0} . Generalizing, if at the time $n \in \mathbb{N}_0$ we have a probability distribution of states p_n on [0, 1], at the time n + 1 the system shall be in the probabilistic state p_{n+1} given by

$$U(p_n) := p_{n+1} = \int_{[0,1]} p_n \left(\delta_{f(y)} \right)_{y \in [0,1]}$$

These considerations allow us to define a probabilistic system generated by a map.

Definition (of probabilistic discrete dynamical system generated by a map). A probabilistic discrete dynamical system on a compact subset K of \mathbb{R}^n generated by a Borelmeasurable function $f : K \to K$ with compact range is defined to be the dynamical system

$$s_f : \mathbb{N}_0 \times P(K) \to P(K)$$

defined recursively by

$$s_f(0,p) = p$$
, $s_f(n+1,p) = \int_K s_f(n,p) (\delta_{f(y)})_{y \in K}$,

for every probabilistic state p and every time n. The evolution operator of s_f is the operator

$$U: P(K) \to P(K): U(p) = \int_{K} p(\delta_{f(y)})_{y \in K}.$$

Remark. Note that the compactness of the range of f guarantees the measurability of δ_f .

Remark. In other terms, the probabilistic discrete dynamical system on a compact subset K of \mathbb{R}^n generated by a Borel-measurable function $f : K \to K$ with compact range is defined to be the dynamical system generated by the function $F : P(K) \to P(K)$ defined by

$$F(p) = \int_{K} p(\delta_{f(y)})_{y \in K}.$$

8. Superpositions of C_c^0 -families in the space $C_c^{0\prime}(T)$

In the following we denote by $C_c^0(T)$ the space of continuous functions with compact support on a locally compact space T, the dual of $C_c^0(T)$ with respect to its standard locally convex topology is the space of measures on T and it's simply denoted by $C_c^{0'}(T)$.

Definition (families of class C_c^0). Let $T \subseteq \mathbb{R}^n$ and $H \subseteq \mathbb{R}^m$ be locally compact subsets. Let $v = (v_i)_{i \in H}$ be a family in the space of measures $C_c^{0'}(T)$ indexed by H. We define image of a continuous test function ϕ with compact support on T by the family v, the function

$$v(\phi): H \to \mathbb{R}: v(\phi)(i) = v_i(\phi).$$

If the image of $v(\phi)$ lies in $C_c^0(H)$, for every $\phi \in C_c^0(T)$, we say that v is a family of class C_c^0 . In this condition we define the operator

$$\widehat{v}: C_c^0(T) \to C_c^0(H): \phi \mapsto v(\phi),$$

and we call it the operator generated by v.

Definition (superposition of measures). In the above conditions, let $a \in C_c^{0'}(H)$, i.e, let a be a measure on H and let v be a family of class C_c^0 in the space $C_c^{0'}(K)$. The functional $a \circ \hat{v}$ is denoted by

$$\int_{H} av,$$

and is called superposition of v by the system of coefficient a. Moreover, if a is a probabilistic system of coefficients, i.e., if a is non-negative and with unitary integral, the superposition $\int_{H} av$ is called a convex superposition of v.

Theorem. Let $a \in C_c^{0'}(H)$, and let v be a family of class C_c^0 in the space $C_c^{0'}(T)$ indexed by the locally compact H. Then the functional

$$\int_{H} av : C_c^0(T) \to \mathbb{R},$$

is a meansure, i.e., a linear continuous functional on $C_c^0(K)$.

Proof. Let K be a compact subset of T, and let ϕ be a real continuous function on T, with compact support contained in K, moreover let H' the support (compact) of $v(\phi)$, we have, for some $j \in H$,

$$\begin{aligned} \left| \left(\int_{H} av \right) (\phi) \right| &= |a \left(v \left(\phi \right) \right)| \le M_{H'}^{a} \| v \left(\phi \right) \|_{C^{0}(H)} = M_{H'}^{a} \max_{i \in H} |v_{i} \left(\phi \right)| = \\ &= M_{H'}^{a} \cdot |v_{j} \left(\phi \right)| \le M_{H'}^{a} \cdot M_{K}^{v_{j}} \cdot \| \phi \|_{C^{0}(K)} \,, \end{aligned}$$

so $\int_{H} av$ is a bounded-linear functional and then it is continuous.

Theorem. Let I, J be intervalls of the real line, $f : I \to J$ be a continuous bijective function and let δ_f be the family $(\delta_{f(x)})_{x \in I}$ in the space $C_c^{0'}(J)$. Then, δ_f is a family of class C_c^0 ; moreover, for every measure $c \in C_c^{0'}(I)$, the superposition $\int_I c\delta_f$ is a measure and we have

$$\left(\int_{I} c\delta_{f}\right)(\phi) = \int_{I} \phi \circ f \ d\mu_{c}.$$

Proof. Note that f is a homeomorphism of I onto J (by a classic result of Analysis). Let us compute $\delta_f(\phi)$ for every $\phi \in C_c^0(J)$; we have

$$\delta_f(\phi): I \to \mathbb{R}, \ \delta_f(\phi)(x) = \delta_{f(x)}(\phi) = \phi(f(x)) = (\phi \circ f)(x),$$

hence $\delta_f(\phi) = \phi \circ f$; note that $\phi \circ f \in C_c^0(I)$, infact,

$$\operatorname{supp}(\phi \circ f) = f^{-}(\operatorname{supp}\phi).$$

Thus δ_f is a C_c^0 -family in $C_c^{0\prime}(J)$. Let $c \in C^{0\prime}(I)$, we have so

$$\left(\int_{I} c\delta_{f}\right)(\phi) = c\left(\phi \circ f\right). \blacksquare$$

9. The generalized Perron-Frobenius operator

In this section we shall prove a generalization of the following result of Prigogine:

Theorem (Prigogine). Let Bernoully shift at time 0 be in a probabilistic-state p_0 , and let p_0 be a regular measure generated by a certain real continuous function g_0 defined on [0, 1]. Then, the orbit $(g_n)_{n \in \mathbb{N}_0}$ starting from g_0 verifies

$$g_{n+1}(x) = \frac{1}{2} \left(g_n\left(\frac{x}{2}\right) + g_n\left(\frac{x+1}{2}\right) \right),$$

for every time $n \in \mathbb{N}_0$.

First, we have to consider a generalization of composition of a measure with a function. Recall that

Definition (of composition in $C_c^{0'}(\Omega)$ **with a diffeomorphism).** Let Ω, Ω' be two open sets of \mathbb{R}^n and let $F \in \text{Diff}^1(\Omega', \Omega)$, that is F is bijective, $F \in C^1(\Omega', \Omega)$, $F^- \in C^1(\Omega, \Omega')$. Then, for every $\psi \in C_c^0(\Omega')$, the function $(\psi \circ F^-) |\det J_{F^-}|$ belongs to the space $C_c^0(\Omega)$, moreover, for every $u \in C_c^{0'}(\Omega)$, the functional

$$\iota \circ F : C_c^0(\Omega') \to \mathbb{K} : \psi \mapsto u\left(\left(\psi \circ F^-\right) |\det J_{F^-}|\right)$$

is a measure on Ω' called the composition of u with F.

In our context we have to define composition in the space $C_c^{0'}(T)$, with T locally compact subspace of \mathbb{R}^n . We recall that T is a locally compact subspace of \mathbb{R}^n if and only if it is the intersection of an open subset with a closed subset of \mathbb{R}^n . The extension is immediate:

Definition (of composition in $C_c^{0'}(T)$ **with a diffeomorphism).** Let T, T' be two locally compact subsets of \mathbb{R}^n , let $F \in \text{Diff}^1(T', T)$, that is there exists a function $F_0 \in \text{Diff}^1(\Omega', \Omega)$, with Ω, Ω' open sets of \mathbb{R}^n containing T and T' respectively, and such that

$$(F_0)_{|(T',T)|} = F.$$

Then, for every $\psi \in C_c^0(T')$, the function $(\psi \circ F^-) |\det J_{F^-}|$ belongs to the space $C_c^0(T)$ and moreover, for every $u \in C_c^{0'}(T)$, the functional

$$u \circ F : C_c^0\left(T'\right) \to \mathbb{K} : \psi \mapsto u\left(\left(\psi \circ F^-\right) |\det J_{F^-}|\right)$$

is a measure on T' called the composition of u with F.

Recall that a real function f, defined on an interval of the real line, is said piecewiseaffine if there is a (ordered) partition $I = (I_j)_{j=1}^k$ of its domain such that each set of I is an interval and f is affine on every set of I. If f_j is the restriction of f to I_j and L_j the slope of f_j , the systems $(f_j)_{j=1}^k$ and $(L_j)_{j=1}^k$ are defined the system of restrictions and the system of slopes of f with respect to I.

Theorem. Let φ_0 be a measure on [0, 1], and let φ be the orbit starting from φ_0 of the probabilistic dynamical system generated by a piecewise-affine mapping $f : [0, 1] \rightarrow [0, 1]$ of partition $(I_j)_{j=1}^k$ and associated system of restrictions $(f_j)_{j=1}^k$. Assume f with compact image (widely) contained in [0, 1] and assume the slopes different from 0. Then, for every time n, we have

$$\varphi_{n+1}\left(\phi\right) = \sum_{j=1}^{k} \frac{1}{|L_j|} \left(\varphi_n \circ f_j^{-}\right) \left(\phi_{|f_j(I_j)}\right),$$

for every $\phi \in C^0([0,1])$, where L_j is the slope of f_j .

Proof. Let $\phi \in C^0([0,1])$, note that the superposition $\int_{[0,1]} \varphi_n \delta_f$ is a measure by a previous result, since f has compact image and it is measurable. Moreover, for each n,

$$\varphi_{n+1}(\phi) = \left(\int_{[0,1]} \varphi_n \delta_f\right)(\phi) = \int_{[0,1]} \phi \circ f \, d\varphi_n = \sum_{j=1}^k \int_{I_j} \phi \circ f \, d\varphi_n = \sum_{j=1}^k \int_{I_j} \phi_{|f_j(I_j)} \circ f_i \, d(\varphi_n)_{|I_j} = \sum_{j=1}^k \left(\int_{I_j} (\varphi_n)_{|I_j} \, \delta_{f_j}\right)(\phi_{|f_j(I_j)}).$$

To justify the last equality note that, for every $\psi \in C_c^0(f_i(I_i))$,

$$\left(\int_{I_j} (\varphi_n)_{|I_j} \,\delta_{f_j}\right)(\psi) = (\varphi_n)_{|I_j} \left(\widehat{\delta}_{f_j}(\psi)\right),$$

in fact, $\delta_{f_i}(\psi)$ is a continuous function with compact support, since, for every y in I_j ,

$$\delta_{f_j}(\psi)(y) = \delta_{f_j(y)}(\psi) = \psi(f_j(y)),$$

hence $\delta_{f_j}(\psi) = \psi \circ f_j$, moreover f_j is a homeomorphism, so $f_j^-(\operatorname{supp}\psi)$ is compact and it is the support of $\psi \circ f_j$. So the family δ_{f_j} is a family of class C_c^0 . Concluding, the superposition

$$\mu := \int_{I_j} \left(\varphi_n \right)_{|I_j} \delta_{f_j}$$

is a measure on I_j . Moreover, since for every $\psi \in C_c^0(f_j(I_j))$ it is

$$\left| \left(\varphi_n \right)_{|I_j} \left(\widehat{\delta}_{f_j} \left(\psi \right) \right) \right| \le \left| \varphi_n \right| \left(\left[0, 1 \right] \right) \left\| \psi \right\|_{C_b^0(I_j)}$$

 μ is a continuous linear functional on $C_c^0(f(I_j))$. Note that, although the function $\phi_{|f_j(I_j)}$ can be not with compact support, it is certainly in $\mathcal{L}^1_\mu(f_j(I_j))$ (it is continuous and bounded on $f_j(I_j)$, i.e., of class C_b^0), and then the expression $\mu(\phi_{|f_j(I_j)})$ is well defined.

Note being the function $f_j : I_j \to f_j(I_j) : x \mapsto L_j x$ an invertible function, by definition of composition of a distribution with a diffeomorphism, we have

$$\left(\int_{I_j} \varphi_n \delta_{f_j}\right)(\psi) = \varphi_n \left(\psi \circ f_j\right) = \frac{1}{|L_j|} \left(\varphi_n \circ f_j^{-1}\right)(\psi),$$

in other words,

$$\int_{I_j} \varphi_n \delta_{f_j} = \frac{1}{|L_j|} (\varphi_n)_{|I_j} \circ f_j^{-1} = \frac{1}{|L_j|} \varphi_n \circ f_j^{-1}$$

and the theorem is completely proved. \blacksquare

We conclude the argumentation generalizing the Perron-Frobenius operator.

Definition (the generalized Perron-Frobenius operator). Let f be a piecewise-affine function on an intervall [a, b] with partition $(I_j)_{j=1}^k$, system of restrictions $(f_j)_{j=1}^k$ and system of slopes $(L_j)_{j=1}^k$. Assume f with compact image (widely) contained in [a, b] and assume the slopes different from 0. We call the operator

$$P: C^{0'}([a,b]) \to C^{0'}([a,b])$$

defined by

$$P(\mu)(\phi) = \sum_{j=1}^{k} \frac{1}{|L_j|} (\mu \circ f_j^{-}) (\phi_{|f_j(I_j)}),$$

for every measure μ on [a, b], the generalized Perron-Frobenius operator associated with the piecewise-affine function f.

References

- [1] I. Antoniou, I. Prigogine, "Intrinsic Irreversibility of Dynamics", Phisica 192 A (1993).
- [2] R. Balescu, Equilibrium and Non-equilibrium Statistical Mechanics (Wiley, New York, 1975).
- [3] I. Prigogine, Non-Equilibrium Statistical Mechanics (Wiley, New York, 1961).
- [4] I. Prigogine, From Being to Becoming (Freeman, New York, 1980).
- [5] I. Prigogine, Le leggi del Caos (Editori Laterza, 1993).
- [6] P. Shields, The Theory of Bernoully Shifts (University of Chicago Press, Chicago, 1973).
- [7] N. Bourbaki, Integration, in four volumes (Hermann, Paris, 1952, 1956, 1959, 1963).
- [8] N. Bourbaki, *Topologie Generale* (Hermann, Paris, 1953, 1955).
- [9] D. Carfi, Teoria delle decisioni (Gabbiano, Messina, 2007).

David Carfi Università degli Studi Messina Facoltà di Economia Via dei Verdi 75 98122 Messina, Italy **E-mail**: davidcarfi@eniware.it

Presented:September 30, 2005Published on line:February 01, 2008